

Prevalence and Determinants of Dating Violence: An Umbrella Review of Systematic Reviews and Meta-analyses

Md Mahbub Hossain¹, Abida Sultana², Qiping Fan², Ping Ma², and Neetu Purohit²

¹Texas A&M School of Public Health

²Affiliation not available

January 02, 2020

Abstract

Objective: Dating violence (DV) is a psychosocial problem which impacts the health and wellbeing of affected individuals. Many studies have reported a high prevalence and various determinants of DV; however, there is a gap of consolidated evidence on this problem. This umbrella review aimed to synthesize the current evidence on the prevalence and determinants of DV from systematic reviews and meta-analyses.

Methods: We systematically searched ten major databases and additional sources to retrieve systematic reviews or meta-analyses reporting prevalence or determinants of DV, published as peer-reviewed journal articles in English language till October 13, 2019. We extracted and synthesized the findings, reported the prevalence of DV, and categorized the determinants of DV using the socioecological model.

Results: We found 16 eligible systematic reviews and meta-analyses. Most of these studies were conducted in developed countries with a majority of adolescent and young adult participants. Studies reported varying prevalence of physical, sexual, psychological/emotional, and technology-assisted DV ranging from 0.1% to 57.5%, 0.1% to 64.6%, 4.2% to 97%, and 5.8% to 92%, respectively. The determinants of DV at different socioecological levels, including individual behavior, substance abuse, psychiatric conditions, experiencing violence and maltreatment; interpersonal factors like family and peer relationships; community and neighborhood characteristics; patriarchy, culture, and socioeconomic inequalities were reported.

Conclusions: These findings suggest a high burden of DV among young populations, who are vulnerable to various socioecological determinants of DV. Future research should examine how these factors influence DV and multi-level interventions should be adopted to address the same.

Prevalence and Determinants of Dating Violence: An Umbrella Review of Systematic Reviews and Meta-analyses

Md Mahbub Hossain*,¹ Abida Sultana,² Qiping Fan,¹ Ping Ma,¹ Neetu Purohit³

Abstract

Objective: Dating violence (DV) is a psychosocial problem which impacts the health and wellbeing of affected individuals. Many studies have reported a high prevalence and various determinants of DV; however, there is a gap of consolidated evidence on this problem. This umbrella review aimed to synthesize the current evidence on the prevalence and determinants of DV from systematic reviews and meta-analyses.

Methods: We systematically searched ten major databases and additional sources to retrieve systematic reviews or meta-analyses reporting prevalence or determinants of DV, published as peer-reviewed journal articles in English language till October 13, 2019. We extracted and synthesized the findings, reported the prevalence of DV, and categorized the determinants of DV using the socioecological model.

Results: We found 16 eligible systematic reviews and meta-analyses. Most of these studies were conducted in developed countries with a majority of adolescent and young adult participants. Studies reported varying prevalence of physical, sexual, psychological/emotional, and technology-assisted DV ranging from 0.1% to 57.5%, 0.1% to 64.6%, 4.2% to 97%, and 5.8% to 92%, respectively. The determinants of DV at different socioecological levels, including

¹ Department of Health Promotion and Community Health Sciences, Texas A&M School of Public Health, College Station, TX 77843, USA. Email: mhossain@tamu.edu

² Nature Study Society of Bangladesh, Khulna 09000, Bangladesh.

³ IIHMR University, Jaipur, Rajasthan 302029, India.

individual behavior, substance abuse, psychiatric conditions, experiencing violence and maltreatment; interpersonal factors like family and peer relationships; community and neighborhood characteristics; patriarchy, culture, and socioeconomic inequalities were reported.

Conclusions: These findings suggest a high burden of DV among young populations, who are vulnerable to various socioecological determinants of DV. Future research should examine how these factors influence DV and multi-level interventions should be adopted to address the same.

Keywords: *Dating Violence; Intimate Partner Violence; Umbrella Review; Systematic Review; Meta-analysis*

Dating violence (DV) is a psychosocial problem affecting the health and wellbeing of many people around the world. The Center for Disease Control and Prevention (CDC) defined DV as a type of intimate partner violence occurring between two people in a close relationship (Center for Disease Control and Prevention (CDC), 2019). Moreover, the United States (US) Department of Justice defined DV as violence committed by an individual who is in a social relationship of a romantic or intimate nature with the victim (The US Department of Justice, n.d.). Similar definitions across different disciplines have conceptualized DV as interpersonal violence including different acts of physical, sexual, or psychological aggression among dating or unmarried romantic partners, who are involved in the mate-selection process (Jouriles, Garrido, Rosenfield, & McDonald, 2009). Physical forms of DV include use of physical forces to cause any form of harms to the target person in a dating relationship whereas sexual DV include physical force to compel a person to engage in a sexual act, or engaging in sexual act in an abusive manner, or involving the victim in a sexual act without willingness or ability to understand the nature of the act (Breiding, Basile, Smith, Black, & Mahendra, 2015). Furthermore, psychological or emotional DV involves acts, threats of acts, or coercive tactics, which may include humiliation, control, isolation, or deliberate actions which may affect the mental state of the victim (Breiding et al., 2015; WHO, 2012).

Different studies have revealed a varying prevalence of DV in different populations. A review of studies conducted in the US has found estimates for dating violence perpetration ranging from 26% to 46% for physical violence and 3% to 12% for sexual violence (Hickman, Jaycox, & Aronoff, 2004). Furthermore, a 10-year (1999–2009) National Teen DV Trend Study found approximately 1 in 10 female adolescents (9.9% to 10.3%) and 1 in 8 male adolescents (9.1% to 11.8%) reported experiencing physical DV in the past year (Howard, Debnam, & Wang, 2013;

Howard, Debnam, Wang, & Gilchrist, n.d.). Moreover, studies from other countries have shown a high burden of DV. A study evaluated DV in samples from the UK and Spain found the prevalence of DV victimization and perpetration as 23% and 30%, respectively (Viejo, Monks, Sánchez, & Ortega-Ruiz, 2016). Another study conducted in Canada found more than 45% of the study participants reported physical DV (Pedersen & Thomas, 1992). A high prevalence of DV may have a wide range of negative health consequences. Victims of DV may have physical trauma and a higher risk of HIV and other sexually transmitted diseases (Teitelman, Ratcliffe, Dichter, & Sullivan, 2008), which may incur high socioeconomic burden. More importantly, DV profoundly impacts the psychosocial processes and mental health of the victims (Temple et al., 2016). Studies have shown high prevalence of psychiatric comorbidities including depression, anxiety, substance abuse, stress disorders, and posttraumatic stress disorder among the victims of DV (Burton, Halpern-Felsher, Rehm, Rankin, & Humphreys, 2016; Duerksen & Woodin, 2019; Y. Park, Mulford, & Blachman-Demner, 2018). The long-term psychosocial outcomes of DV may include fear to new relationship, difficulties in establishing meaningful communication, social isolation, loneliness, and increased risks of subsequent DV victimization and perpetration (Duerksen & Woodin, 2019; Y. Park et al., 2018).

To address the epidemiological and psychosocial challenges related to DV, it is essential to understand the magnitude of the condition across different population groups. It may help in identifying the high-risk groups and adopt preventive and therapeutic measures to alleviate the burden of DV. Moreover, intrapersonal characteristics, interpersonal relationship dynamics, and sociodemographic factors measures play a critical role in determining the burden of DV in any given population (Foshee et al., 2008). Therefore, it is necessary to examine the determinants of DV to better understand the underlying factors associated with the burden of DV across diverse

population groups. Several reviews exist which reported either the prevalence or factors associated with DV, but there is no review of the systematic reviews and meta-analyses offering consolidated evidence on prevalence and determinants of DV. Considering there are about 11 systematic reviews published each day, there is a critical need for synthesizing evidence from existing reviews and meta-research to advance knowledge and practice (Aromataris et al., 2015). This umbrella review aims to synthesize the current evidence on the global prevalence and determinants of DV from systematic reviews and meta-analyses, which may inform future scientific, therapeutic, and policy discourses in DV.

Methods

Searching the Literature

To conduct this umbrella review, we used the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (Liberati et al., 2009) and the Joanna Briggs Institute (JBI) methodology for umbrella reviews (Aromataris et al., 2015). We searched MEDLINE, EMBASE, PubMed, Academic Search Ultimate, CINAHL, PsycINFO, Violence and Abuse Abstracts, Health Source Nursing/Academic Edition, Health Policy Reference Center, and the Cochrane Library to retrieve potential studies relevant to the objective of this umbrella review. To identify systematic reviews and meta-analyses on prevalence and correlates of DV, we used the following search syntax: (“date violence” OR “date abuse” OR “date aggression” OR “dating violence” OR “dating abuse” OR “dating aggression” OR “courtship violence” OR “courtship abuse” OR “premarital abuse” OR “premarital violence” OR “premarital aggression”) AND (prevalence OR incidence OR statistics OR epidemiology OR predictors OR correlates OR determinants OR "risk factors" OR "associated factors" OR "factors associated" OR factors) AND ("systematic review" OR "meta-analysis" OR "pooled estimate" OR "pooled prevalence")

OR "pooled effect"). The keywords were used to search titles, abstracts, subject headings (like MeSH), and general search terms. No language restriction was applied during database searching. All the databases were searched since the inception of the respective databases. The first search was conducted on August 27, 2019 and same search process was repeated on October 13, 2019, for the last time. In addition to database searching, we retrieved additional articles through hand-searching violence and psychiatry-related journals and communicating with subject-matter experts. Moreover, we used the "cited by" function in Google Scholar to check the eligibility of the citing articles of the primarily screened articles. All the retrieved citations were uploaded to RefWorks and Rayyan software for citation management and systematic evaluation processes. After removing the duplicates, two authors independently evaluated the titles and abstracts of the citations as per the selection criteria of our review. A third author was consulted after the primary screening to confirm the eligibility of the included articles, resolve the conflicts in the selection process, and reach a consensus about the finally recruited articles.

Selection Criteria

In this umbrella review, we included an article if it: a) was a systematically conducted literature review or research synthesis using qualitative (meta-synthesis) or quantitative (meta-analysis) approaches, b) primarily focused on any type of DV among partners in a dating or romantic relationship, c) synthesized either the prevalence or correlates of DV from the primary studies in respective reviews, d) included full-text in English, e) published as a journal article.

We excluded an article if it: a) was not a systematically conducted review (for example, non-systematic literature reviews, overviews, primary articles, opinions, letters, editorials were excluded), b) did not primarily focus on any form of DV in dating relationships (for example, reviews focusing on domestic or any other violence, or articles presenting dating violence with

other types of violence together were excluded), c) did not report either prevalence (including incidence or proportion in a given population, if reported) or determinants (also can be termed as correlates or associated factors in different methods and study designs) of DV synthesized from primary articles, d) did not have the full-text in English, e) was not a journal article (for example, dissertations, policy briefs, reports, book chapters were excluded).

Evaluation of the Methodological Quality of the Recruited Studies

Critical appraisal of the methodological quality of the recruited reviews is an essential process of conducting an umbrella review (Aromataris et al., 2015). In this review, we evaluated the quality of the systematic reviews and meta-analyses using the JBI critical appraisal checklist for systematic reviews and research synthesis checklist. This checklist facilitates distinguishing the quality of studies and decision-making about the recruitment of the articles. Moreover, this checklist has ten items; each item can receive one point and the overall quality score of a study can range from zero to ten. In this umbrella review, studies receiving zero to four, five to seven, and eight to ten were categorized as low, medium, and high-quality studies, respectively.




Figure 1: Flow diagram of the review

Data Extraction and Synthesis

A codebook for extracting quantitative and qualitative data was prepared using the JBI data extraction tool for systematic reviews and research synthesis (Munn, Tufanaru, & Aromataris, 2014). Details on the authors, year of publication, characteristics of the review studies, origin and types of the primary studies in each review, sample size and characteristics of the study population, and key findings on prevalence, determinants, correlates, or associated factors of DV were extracted by two authors and reviewed by a third author. The extracted data were narratively synthesized to present the characteristics of the review studies and the findings meeting the objective of this study. Prevalence data were presented as pooled estimates from meta-analysis, as reported in narrative reviews. Moreover, determinants, correlates or factors associated with DV were extracted and evaluated using socioecological model of McLeroy and colleagues (McLeroy, Bibeau, Steckler, & Glanz, 1988). Using this approach, different variables associated with DV were categorized as individual, interpersonal, community, social, and systems-level determinants of DV. Qualitative or narrative findings were presented as they were reported in the respective reviews, whereas quantitative data were presented with the range of rates within 95% confidence interval (CI), wherever reported.

Results

We found 1119 citations from database searching and 416 citations from additional sources, totaling 1535 citations. After removing the duplicates, screening the titles/abstracts, and evaluating the full texts, we found 16 articles that met our criteria (Caridade, Braga, & Borrajo, 2019; Duval, Lanning, & Patterson, 2018; Garthe, Sullivan, & McDaniel, 2017; Hébert et al., 2019; Johnson et al., 2017; Johnson, Parker, Rinehart, Nail, & Rothman, 2015; Joly & Connolly, 2016; S. Park & Kim, 2019, 2018; Rothman, McNaughton Reyes, Johnson, & LaValley, 2012;

Rubio-Garay, López-González, Carrasco, & Amor, 2017; Spencer, Toews, Anders, & Emanuels, 2019; Stonard, Bowen, Lawrence, & Price, 2014; Taquette & Maia Monteiro, 2019; Wincentak, Connolly, & Card, 2017; Zych, Viejo, Vila, & Farrington, 2019). A flow-diagram of the screening process is depicted in Figure 1.

Characteristics of the Studies and Participants

Among the 16 studies included in this review (summary of the studies is provided in Table 1), ten were meta-analyses (Garthe et al., 2017; Hébert et al., 2019; Johnson et al., 2017; Joly & Connolly, 2016; S. Park & Kim, 2018, 2019; Rothman et al., 2012; Spencer et al., 2019; Wincentak et al., 2017; Zych et al., 2019) and six were systematic reviews (Caridade et al., 2019; Duval et al., 2018; Johnson et al., 2015; Rubio-Garay et al., 2017; Stonard et al., 2014; Taquette & Maia Monteiro, 2019). These studies were published from 2012 to 2019, while most (n = 15) of the studies were published since 2017. The number of primary studies within those reviews ranged from 13 to 113, and the sample sizes ranged from 24 to 85,198. While most participants across all the studies were adolescents, four reviews (Spencer et al., 2019; Taquette & Maia Monteiro, 2019; Wincentak et al., 2017; Zych et al., 2019) exclusively recruited primary studies with adolescent participants, whereas remaining reviews recruited studies comprising adolescents, young adults, and participants of other age groups. Most primary studies in all the reviews were conducted in the US and Canada; however, four reviews included primary studies conducted in different states of the US only (Duval et al., 2018; Johnson et al., 2017, 2015; Spencer et al., 2019). Very few primary studies were reported from low- and middle-income countries.

Table 1: Characteristics and the key findings of the included systematic reviews and meta-analyses

Source	Name and timeframe of searching Databases	Number and type of primary studies	Origin of included studies	Sample size and characteristics of the study population (if reported)	Key findings related to prevalence and/or associated factors of DV
(Rothman et al., 2012)	PsycINFO, ISI Web of Knowledge (including citations from Medline, BIOSIS, and other databases); 1985-2010	28 (82% cross-sectional)	The US (n = 23), Canada (n = 1), Mexico (n = 1), Russia (n = 1), New Zealand (n = 1), South Africa (n = 1)	Sample size ranged from 40 to 13,422; age ranged from 12 to 54 years; 7 studies had high school participants, 14 had college or university-based samples, 7 had participants from communities and other settings	Odds ratio for DV perpetration for heavy episodic drinking, problem use, and frequency/quantity were 1.47 (95% CI: 1.17, 1.85), 2.33 (95% CI: 1.94, 2.8), and 1.23 (95% CI: 1.16, 1.31) in fixed-effect model
(Stonard et al., 2014)	Academic Search Complete, PsycINFO, Science Direct, Google Scholar; Post-2000 studies	65 (48 cross-sectional, 9 longitudinal, and 8 retrospective studies)	United States (n = 19), UK (n = 17), Canada (n = 16), Germany (n = 4) Sweden (n = 3), Spain (n = 2), New Zealand (n = 2), Netherlands (n = 1), Switzerland (n = 1)	Sample size ranged from 117 to 81247; age ranged from 10 to 29 years; most studies recruited adolescent participants	12% to 56% participants reported some form of victimization, 12% to 54% reported perpetration of Technology Assisted Adolescent Dating Violence and Abuse (TAADVA). Physical violence victimization (ranged from 10% to 30%), physical violence perpetration (5% to 30%), psychological or emotional violence victimization (35% to 55%), psychological or emotional violence perpetration (20% to 70%), sexual violence victimization (5% to 30%), and sexual violence perpetration (5% to 20%) were reported

(Johnson et al., 2015)	PubMed, Web of Science, PsycINFO; 2005-2015	20 (14 cross-sectional studies)	Different states of the US	11 studies had male and female participants, 9 had youth from both urban and non-urban settings, 6 focused on adolescents, 9 focused on emerging adults; 15 of the 20 studies recruited participants from school settings	Neighborhood factors associated with DV included social disorganization (8 studies evaluated association), five studies reports association with collective efficacy, neighborhood disorder (1 out of 12 studies reported association with DV), alcohol outlets (4 studies reported association with DV), demographic and structural challenges (11 studies assessed neighborhood-level structural factors, 8 investigated association of residence characteristics with DV). Neighborhood disorder is associated with physical dating violence perpetration, but do not with physical dating violence victimization.
(Joly & Connolly, 2016)	PsycINFO, ERIC, and Social Sciences Abstracts; timeframe not specified	29 (21 quantitative and 8 qualitative studies)	Most studies were from the US; 1 qualitative study each from Canada and UK	Sample size ranged from 24 to 724 (for quantitative studies) and 1 to 29 (for qualitative studies); high-risk population were surveyed including street-involved, pregnant, maltreated, juvenile justice system participants	34% (95% CI 24% to 45%) participants experienced physical DV, 45% (95% CI 31% to 61%) participants perpetuated physical DV; factors influencing DV included vulnerability due to being high-risk individuals, teenage mothers, juvenile justice system, truancy, psychological conditions like hallucination, delusion, drug use, self-harm, suicidal ideation, irrational thinking, and post-traumatic stress disorder; social factors associated with DV included negative peer relations, lack of family and social support, negative neighborhood experience.

(Rubio-Garay et al., 2017)	MEDLINE, PsycINFO, Academic Search Premier, SCOPUS, Web of Science, and additional sources like e-journals; studies published till 2013	113 (all quantitative studies)	Most studies were conducted in the US and other developed countries	Sample size ranged from 490 to 81247; most studies included adolescents followed by young adult and mixed populations	The prevalence of different forms of DV was reported as: perpetrated physical violence (3.8% to 41.9%), victimized physical violence (0.4% to 57.3%), perpetrated psychological violence (4.2% to 97%), victimized psychological violence (8.5% to 95.5%), perpetrated sexual violence (1.2% to 58.8%), victimized sexual violence (0.1% to 64.6%).
(Garthe et al., 2017)	PsycINFO, Web of Science, and relevant journals; 2000-2014	27 (all quantitative studies)	One study each was conducted in Brazil and Thailand; all the remaining were from the US (n = 15) and Canada (n = 10)	Sample ranged from 43 to 4131; all samples included adolescent participants	Three peer risk factors of DV were meta-analyzed; peer dating violence ($r = .30$), peers' aggressive and/or antisocial behavior ($r = .20$), and being victimized by peers ($r = .22$) were all significantly related to DV perpetration and victimization among the adolescents.
(Johnson et al., 2017)	MEDLINE and PsycINFO; 2003-2015	13 (7 cross-sectional, 5 longitudinal, 1 retrospective)	Different states of the US	Sample size ranged from 67 to 15123; participants were adolescents and emerging adults	Marijuana use was associated with physical DV victimization (OR 1.54, 95% CI: 1.22-1.93) and perpetration (OR 1.45, 95% CI: 1.2-1.75)
(Wincentak et al., 2017)	PsycINFO, Sociological Abstracts, and the Education Resources	101 (all quantitative)	Not specified	Sample size ranged from 48 to 81247; participants aged 13 to 18 years	The overall prevalence of physical and sexual DV was 20% (95% CI: 17%-23%) and 9% (95% CI: 5%-14%) respectively with significant variability across studies; physical DV ranged from 1% to 61%, and sexual DV ranged from less than 1% to 54%. Boys and girls

	Information Center (ERIC); 1980-2013				experienced similar rate (21%) of victimization; however, boys experienced less (13%) perpetration compared to girls (25%). Girls reported lower rates of perpetration compared with boys (3% vs 10%) and higher rates of victimization (14% vs 8%). Sample groups with higher prevalence had older teens (sexual DV), cultural minority girls (physical DV), and disadvantaged neighborhoods (physical DV)
(Duval et al., 2018)	EBSCO, Scopus, PsycINFO, MEDLINE, ERIC, and PubMed; 2006-2016	23 (22 quantitative, 1 mixed method)	Different states of the US	Sample size ranged from 43 to 5035, aged between 18 to 22.06 years; undergraduate students from different academic institutions	Female participants were more likely to perpetrate and/or experience DV than males. Up to 43.3% males perpetrated DV whereas up to 50.4% females were victims of DV. Individual risk factors included substance use, high-risk sexual behaviors; family risk factors included intergenerational violence; and peer and social risk factors included Greeklife or athletic team membership. Self-control and positive relationship type were reported as protective factors.
(S. Park & Kim, 2018)	PubMed, Web of Science, and SCOPUS; studies published till 2016	27 (all quantitative studies)	The US (n = 19), Canada (n = 6), Taiwan (n = 1), Switzerland (n = 1)	15 studies included teenagers and 12 studies included adults; age ranged from 11 to 21.5	Correlates of DV included family related factors like family structure and relationship, parenting, witnessing IPV, childhood maltreatment, fear of family violence, parents' education, socioeconomic status. Community related correlates are bullying, deviant peers, positive friendship, grade, school attachment, neighborhood hazard and support. Having deviant peers was the

					strongest risk factor of DV perpetration, whereas witnessing parental violence was the strongest risk factor of DV victimization. Also, risk factors of DV were more powerful predictors of DV perpetrators and victims than the protective factors.
(Caridade et al., 2019)	Academic Research Complete, Business Source Complete, Complementary Index, ERIC/EBSCOhost, Psychology and Behavioral Sciences Collection, PubMed, Science Direct, SCOPUS, and Social Sciences Citation Index; studies published till 2018	44 (most studies were cross-sectional)	Spain (n = 8), Mexico (n = 3), one study each from Belgium, Israel, Italy, Canada, UK, and Portugal; remaining studies from the US	Sample size ranged from 155 to 5647, age ranged from 12 to 25 years; most studies recruited students from academic and community settings	Prevalence of cyber DV perpetration ranged from 8.1% to 93.7%; prevalence of DV victimization ranged from 5.8% to 92%; cyber DA perpetration was related to legitimization of DV by boys, behavioral or romantic jealousy, beliefs in myths of love, sexist beliefs, endorsement of gender stereotypes, narcissism vulnerability and grandiosity, experiencing other forms of violence, offline DV experience, family violence, bullying and cyber bullying; cyber victimization was related to delinquency or antisocial behaviors, other high-risk behavior, initiated sexual activity, time spent on social networking, higher anxiety or depressive symptoms, other psychological issues, sexual orientation, worse dyadic adjustment.
(Hébert et al., 2019)	PsycINFO, PubMed, Google Scholar, and relevant French	87; study types were not reported	The US (72%), Canada (19%), other studies from Asia (South Korea	76 studies included in meta-analysis had a total sample was 278,712 participants;	Peer risk factors included peer victimization ($r = .186$), peer sexual harassment ($r = .297$), deviant peers ($r = .25$); protective factors included parental monitoring ($r = -.126$), parental support ($r = -.109$), peer support ($r = -.139$)

	databases; 2000-2015		and India), Europe (Spain), and South America (Mexico, Chile, and El Salvador)	sample ranged from 41 to 85,198; 62% samples included adolescents and 33% had emerging adult participants	
(S. Park & Kim, 2019)	PubMed, Web of Science, and SCOPUS; studies published till 2016	25 (all quantitative studies)	Place not specified	Mean age of the participants ranged from 13.8 to 23; most participants were adolescents or young adults	DV victims and perpetrators were more likely to be involved in violence experiences. DV victims or DV perpetrators were more likely to assume the opposite role as well. Also, DV perpetrators had a strong association with previous or concurrent experience of DV victimization, and current DV victims had experiences of being both victim and perpetrator. DV victimization had a stronger association with general violence experiences than did DV perpetration
(Spencer et al., 2019)	PsycINFO, PubMed, Proquest, Proquest Dissertations and Theses, Education Resources Information Center (ERIC), and Social Services	37 (20 longitudinal, 17 cross-sectional)	Different states of the US	Sample size included in the meta-analysis was 33,537; all studies recruited adolescent participants	Physical DV victimization was the strongest risk marker for physical DV perpetration. Also, externalizing behaviors, anger, witnessing violence, approval of violence, risky sexual behaviors, controlling behavior, substance abuse, depression, childhood maltreatment and delinquency were the strongest risk markers for DV perpetration. In contrast, conflict resolution skills, relationship quality with parents, and responsibility were protective markers against TDV

	Abstracts; 1997-2018				perpetration. DV perpetration and depression were significantly stronger risk markers for female perpetration whereas controlling behaviors was a significantly stronger risk marker for male TDV perpetration
(Taquette & Maia Monteiro, 2019)	PubMed; studies published till 2017	35 (25 cross-sectional, 9 longitudinal, 1 qualitative)	The US (n = 25), Spain (n = 2), South Africa (n = 2), and one study each in Canada, Egypt, Liberia, Malawi, Rwanda, Zambia, Sweden, and the UK	Sample size ranged from 39 to 75590; all participants were adolescents	Varying prevalence of DV was reported across studies (up to 70.7%), factors associated with DV included patriarchal culture, racism, heterosexism, poverty, contextual issues in family, school, community and social media, health problems including depression, anxiety, low self-esteem, substance abuse, and high-risk sexual behavior
(Zych et al., 2019)	Web of Science, Google Scholar, SCOPUS; studies published till 2016	23 (all quantitative studies)	The US (n = 15), Canada (n = 6), one study each from Norway and South Africa	Sample size ranged from 88 to 17,780; all participants were adolescents	DV perpetration was associated with bullying perpetration (AOR = 1.29, 95% CI: 1.07-1.56), DV victimization was associated with bullying perpetration (AOR = 1.09, 95% CI: 1.01-1.12), which was stronger for males. Bullying victimization was related to DV victimization (AOR = 1.96, 95% CI: 1.68-2.29), which was stronger for females. The relation between bullying victimization and DV perpetration was not statistically significant (AOR = 1.01, 95% CI: 0.99-1.03)

For example, among 87 studies included in a meta-analysis by Hébert and colleagues, only two primary studies were from India and El Salvador (Hébert et al., 2019). Most of the studies (n = 11) (Duval et al., 2018; Garthe et al., 2017; Hébert et al., 2019; Johnson et al., 2017; S. Park & Kim, 2018, 2019; Rothman et al., 2012; Spencer et al., 2019; Stonard et al., 2014; Wincentak et al., 2017; Zych et al., 2019) included in this umbrella review had high quality whereas remaining (n = 5) (Caridade et al., 2019; Johnson et al., 2015; Joly & Connolly, 2016; Rubio-Garay et al., 2017; Taquette & Maia Monteiro, 2019) studies had medium quality as per the JBI quality appraisal checklist (Appendix 1).

Prevalence of DV

Physical DV. Studies have reported a varying prevalence of physical DV. Stonard and colleagues reported the prevalence ranging from 5% to 30% and 10% to 30% for physical DV perpetration and victimization, respectively (Stonard et al., 2014). Another systematic review by Rubio-Garay and colleagues reported the rates of physical DV perpetration and victimization, ranging from 3.8% to 41.9% and 0.4% to 57.5%, respectively (Rubio-Garay et al., 2017). Two meta-analyses reported the pooled prevalence of Physical DV (Joly & Connolly, 2016; Wincentak et al., 2017). In the meta-analysis by Joly and colleagues, 34% (95% CI: 24%-45%) participants experienced physical DV, whereas 45% (95% CI: 31%-61%) participants perpetrated physical violence in their dating relationships (Joly & Connolly, 2016). Another meta-analysis of 96 studies found the overall prevalence of reported physical DV was 20% (95% CI: 17%-23%) (Wincentak et al., 2017).

Sexual DV. A meta-analysis by Wincentak and colleagues found the prevalence of sexual DV as 9% (95% CI: 5%-14%) (Wincentak et al., 2017). Moreover, Duval and colleagues reported 43.3% of men claimed to perpetrate sexual DV, whereas 50.4% of women reported being a

victim of sexual DV (Duval et al., 2018). Stonard and colleagues found the rates of sexual DV perpetration and victimization ranging from 5% to 20% and 5% to 30% respectively (Stonard et al., 2014). Furthermore, Rubio-Garay and colleagues reported 1.2% to 58.8% of participants perpetrated sexual DV whereas 0.1% to 64.6% reported being victims of sexual DV (Rubio-Garay et al., 2017).

Psychological and Emotional DV. Studies have also evaluated the prevalence of psychological and emotional DV (Rubio-Garay et al., 2017; Stonard et al., 2014). Stonard and colleagues found 35% to 55% of participants reported psychological or emotional DV victimization whereas 20% to 70% reported perpetration of such violence (Stonard et al., 2014). Another review by Rubio-Garay and colleagues revealed the prevalence of psychological DV victimization and perpetration ranging from 8.5% to 95.5% and 4.2% to 97% respectively (Rubio-Garay et al., 2017).

Other Forms of DV. In addition to the above-mentioned typologies, DV is also reported to happen in digital platforms. Stonard and colleagues found the prevalence of technology-assisted DV perpetration ranged from 12% to 54% whereas victimization ranged from 12% to 56% among the study participants (Stonard et al., 2014). DV in the digital era was also evaluated by Caridade and colleagues who found the prevalence of cyber dating abuse perpetration ranging from 8.1% to 93.7% and victimization ranging from 5.8% to 92% (Caridade et al., 2019).

Determinants of DV

Individual-level determinants. Several determinants of DV were reported across studies that may influence the perpetration or victimization of DV among the individuals. Personality and behavioral attributes like low esteem (Taquette & Maia Monteiro, 2019), bullying (Caridade et

al., 2019; Zych et al., 2019), narcissism vulnerability (Caridade et al., 2019), grandiosity (Caridade et al., 2019), and controlling behavior (Spencer et al., 2019) were found to be associated with DV. Moreover, alcoholism and other forms of substance abuse were associated with DV (Caridade et al., 2019; Duval et al., 2018; Johnson et al., 2017, 2015; Joly & Connolly, 2016; Rothman et al., 2012; Spencer et al., 2019; Taquette & Maia Monteiro, 2019). For example, a meta-analysis by Rothman and colleagues reported higher odds ratio (OR) of DV perpetration for heavy episodic drinking (OR 1.47, 95%CI: 1.17-1.85), problematic use (OR 2.33, 95%CI: 1.94-2.8), and frequency/quantity of drinking (OR 1.23, 95%CI: 1.16-1.31) (Rothman et al., 2012). Furthermore, psychiatric disorders like depression, suicidal ideation, and posttraumatic stress disorder were associated with DV perpetration and victimization (Caridade et al., 2019; Joly & Connolly, 2016; Spencer et al., 2019; Taquette & Maia Monteiro, 2019). In addition, poor educational attainment (S. Park & Kim, 2018), high-risk sexual behavior (Caridade et al., 2019; Duval et al., 2018; Spencer et al., 2019; Taquette & Maia Monteiro, 2019), and belonging to vulnerable groups like teenage mothers (Joly & Connolly, 2016) were associated with DV. Also, previous exposure to any forms of violence, including childhood abuse or domestic violence, was associated with DV perpetration and victimization (Caridade et al., 2019; Duval et al., 2018; S. Park & Kim, 2018, 2019; Spencer et al., 2019; Zych et al., 2019). For example, a meta-analysis by Park and Kim revealed that participants who experienced violence earlier were more likely to perpetrate or become the victims of DV (S. Park & Kim, 2019). While most of the determinants were reported as risk factors for DV, individual characteristics like self-control (Hébert et al., 2019) and conflict resolution skills (Spencer et al., 2019) have been reported as protective factors against DV.

Interpersonal determinants. Interpersonal relationships and interactions have been considered as critical elements in DV studies. The characteristics of the peers and the quality of interactions with them are found as major determinants of DV (Duval et al., 2018; Garthe et al., 2017; Hébert et al., 2019; Joly & Connolly, 2016; S. Park & Kim, 2018, 2019; Spencer et al., 2019; Zych et al., 2019). For example, a meta-analysis by Garthe and colleagues reported peers' aggressive or antisocial behavior ($r = .20$), peer DV ($r = .30$), and being victimized by peers ($r = .22$) were significantly associated with DV perpetration and victimization (Garthe et al., 2017). Moreover, family structure and relationships were associated with DV among children, adolescents, and young adults (Joly & Connolly, 2016; S. Park & Kim, 2018; Spencer et al., 2019; Taquette & Maia Monteiro, 2019). For example, the meta-analysis by Park and Kim found that several family factors predicted DV, among which “witnessing parental violence” was the strongest risk factor of DV victimization (S. Park & Kim, 2018). In contrast, parent support (Hébert et al., 2019) and relationship quality with parents (Spencer et al., 2019) were protective factors against DV.

Community-level level determinants. Studies found that community and neighborhood characteristics often influence DV perpetration and victimization (Johnson et al., 2015; Joly & Connolly, 2016; Wincentak et al., 2017). Johnson and colleagues found neighborhood disorder, disorganization, demographic and structural challenges, and residence characteristics were associated with DV perpetration but not with victimization (Johnson et al., 2015). A meta-analysis Joly and colleagues (Joly & Connolly, 2016) reported negative neighborhood experience was associated with DV whereas samples from disadvantaged neighborhoods had higher physical DV is a meta-analysis conducted by Wincentak and colleagues (Wincentak et al., 2017).

Social and systems-level determinants. Several social and systems-level determinants of DV were reported across studies. Social norms like patriarchal social structure (Taquette & Maia Monteiro, 2019) and legitimacy of DV by boys (Caridade et al., 2019) were associated with DV. Moreover, social structures, institutions, and practices were associated with power imbalance and vulnerability to DV (Johnson et al., 2015; Joly & Connolly, 2016; Taquette & Maia Monteiro, 2019; Wincentak et al., 2017). For example, moderator analyses in the meta-analysis by Wincentak and colleagues found samples with socioeconomically disadvantaged and cultural minority girls had higher rates of physical DV (Wincentak et al., 2017). Moreover, Joly and colleagues found participants in the juvenile justice system were more vulnerable to DV (Joly & Connolly, 2016). Furthermore, Taquette and colleagues reported social and systems-level determinants like culture, racism, sexism, and poverty were associated with DV perpetration and victimization (Taquette & Maia Monteiro, 2019).

Discussion

This umbrella review evaluated the current evidence on the prevalence and determinants of DV from published systematic reviews and meta-analyses. The findings suggest a high burden of different forms of DV perpetration and victimization across different population groups. In addition, the determinants of DV at different levels of the socioecological model inform how individual, interpersonal, community, social and systems-level factors may influence DV among the study participants. These findings offer several critical insights about DV, which are essential for future research, policy development, and practice.

First, the reported prevalence of DV may provide a sub-optimal picture of the actual burden. A possible reason for this may be under-reporting due to stigma and other psychosocial barriers perceived by the affected individuals (Foshee et al., 2008; Hickman et al., 2004; Temple et al.,

2016). Healthcare professionals, social workers, legal service providers, and other stakeholders in health and social care should provide support and create an enabling environment that helps the victims to report the cases of DV (Hickman et al., 2004; Y. Park et al., 2018). Another reason may be attributable to the definitions and constructs of DV across studies published from different disciplines. Differences in operational definitions of DV and methods to measure the same may contribute to heterogeneity in estimating the problems and factors associated with DV (Stonard et al., 2014). Future scientific discourses should standardize the definitions and methods, which may facilitate more accurate estimations of the prevalence and determinants of DV.

Second, most of the study participants in the primary studies of the included reviews were adolescents and young adults. A high prevalence of DV among this young population illustrates profound mental health implications. Moreover, DV was found to be associated with behavioral abnormalities, alcoholism and substance abuse, and several psychiatric comorbidities (Burton et al., 2016; Duerksen & Woodin, 2019; Y. Park et al., 2018; Taquette & Maia Monteiro, 2019). This may create a complex scenario of psychiatric multimorbidity among individuals who perpetrate or experience DV. Identifying such cases and addressing the same with multipronged mental health services can be difficult. Mental health researchers and practitioners may consider future research and collaborative actions to better understand such psychosocial complexities associated with DV, which may help in understanding and managing such problems.

Third, most primary studies in this review were conducted in the US, Canada, and countries with similar socioeconomic contexts. A lack of published studies from resource-constrained countries indicates substantial gaps in the research activities or publications in this domain. This appears to be paradoxical as many low- and middle-income countries have a high burden of child marriage

and gender-based violence. It is difficult to say whether the concept of dating violence is not yet established in those contexts, or the dominant paradigm of violence research is concentrated on gender-based violence in those countries. However, very few studies from developing countries have shown a similar burden of DV in those contexts highlighting prospects for future research. Moreover, with demographic and epidemiological transitions in those countries, many people are having increased life expectancy and getting married at a later age, which is contributing to a reduction in child marriage in many countries (Das Gupta et al., 2014). Therefore, a larger proportion of the adolescent and young adult population who are unmarried, may get involved in dating relationships. Given the existence of the determinants of DV revealed in this review, those individuals may experience DV in their dating relationships, which is essential to be measured, prevented, and treated, whenever it occurs.

Fourth, the emergence of technology-assisted or cyber DV in recent years is a major mental health concern. With the rapidly increasing number of people using gadgets and having access to the internet may contribute to the growing burden of such violence among people who share the psycho-epidemiological characteristics of DV perpetration or victimization. The similarities and differences of such cases with other forms of DV may help in understanding the phenomenology of DV in the era of digital technologies and address the same. Moreover, higher access to technologies also offers unique opportunities to use such platforms for delivering interventions promoting awareness and preventing DV (Draucker et al., 2019; O'Brien, Sauber, Kearney, Venaglia, & Lemay, 2019), which should be evaluated before large-scale adoption of such approaches.

Lastly, different determinants of DV at multiple levels of the socioecological model inform the complexities among the levels and how they affect DV among the individuals. Risk factors

identified at the individual level can help in improving the existing guidelines and approaches to diagnose or predict the potential cases of DV. Moreover, individual characteristics may be helpful in designing more personalized care for DV-affected people. Also, interpersonal factors like peer and family relationships highlight the role of dyadic interactions on the psychosocial wellbeing of the individuals. It is essential to consider critical factors like deviant peers or negative family relationships in DV research and practice to alleviate the burden of DV. In addition, community-level characteristics like neighborhood problems may necessitate a multi-sectoral and participatory approach involving community members, planners, social workers, development organizations, and other stakeholders to identify the prevalent causes of DV in the community settings. Furthermore, social and systems-level determinants inform the macro-environmental attributes of DV, which should be addressed in a collective manner. For example, educational institutions are part of the social system, which may play critical roles in delivering interventions preventing DV (Carlos et al., 2017). Such opportunities should be explored, and future interventions should incorporate behavior change constructs considering diverse social and institutional factors, which may increase awareness about DV and prevent the same at the population-level. Nevertheless, social challenges including racism, sexism, impaired health rights in juvenile justice systems, and structural inequalities should be carefully examined and addressed at policy level to prevent the DV epidemic, as those issues continue to affect the mental state of people.

This review has several limitations, which should be acknowledged. Although we searched major databases, but there may be additional literature which we could not evaluate in this review. Another limitation can be the publication bias. Systematic reviews and meta-analyses with insignificant findings may not have been published, which remain beyond the scope of our

review. Moreover, we included studies that explicitly focused on DV. Studies focusing on domestic or gender-based or intimate partner violence were excluded from this review. Those studies may have included DV-related information, which may have provided different insights into this area. Furthermore, we categorized the determinants using the socioecological model. In this process, one determinant may have appeared at multiple levels. For example, previous experience of violence was reported at the individual level, whereas violence at peer or family relationships was reported at the interpersonal level. Such overarching and interconnected factors should be deeply investigated in case of DV to address the problems. Lastly, we did not compare individual primary studies and kept the scope of our review within the reviews and meta-analyses. An in-depth and different organization of individual studies may have considered the in-between studies offering different understanding than an umbrella review approach. However, we conducted this review using state-of-the-art guidelines, and each step of the review process was conducted by more than two authors to minimize biases. Future research should address our limitations and advance the evidence base on DV.

Conclusion

This umbrella review synthesized the current evidence on the prevalence and determinants of DV from systematic reviews and meta-analyses. This evidence provides an overall understanding of DV among the study populations. A high prevalence and various socioecological determinants of DV may inform the development and implementation of policies and interventions to address DV across populations. Also, the research gaps found in the existing literature may inform prospective measures to advance the existing knowledge base. The collective insights of this umbrella review should be used to promote better research and evidence-based approaches preventing DV among people who are or will be in dating relationships.

Critical findings of this umbrella review:

1. This review identified 16 eligible systematic reviews and meta-analyses reporting the prevalence and determinant of DV.
2. In the included reviews, most of the primary studies were conducted in developed countries, and the majority of the participants were adolescents and young adults
3. Studies reported varying prevalence of physical, sexual, psychological/emotional, and technology-assisted DV ranging from 0.1% to 57.5%, 0.1% to 64.6%, 4.2% to 97%, and 5.8% to 92%, respectively.
4. The socioecological determinants of DV included individual behavior, substance abuse, psychiatric conditions, experiencing violence and maltreatment; interpersonal factors like family and peer relationships; community and neighborhood characteristics; patriarchy, culture, and socioeconomic equalities.

Implications on Practice, Research, and Policy

To the best of our knowledge, this umbrella review is the first one to examine the consolidated evidence on the prevalence and determinants of DV. This review found a high prevalence and diverse socioecological determinants of DV, which may have following implications for practice, research, and policy:

1. Evidence synthesized in this review provide insights on the severity and determinants of DV, which inform how different populations at-risk should be targeted for evidence-based preventive measures.

2. An understanding of socioecological determinants may allow the development and implementation of interventions improving behavioral and psychiatric conditions of the individuals, and enhancing the nature and quality of family and peer relationships
3. Policies and programs should be adopted to improve the neighborhood characteristics, address cultural disparities, and alleviate social inequalities, which may contribute to a long-term prevention of DV in a given population.
4. The gaps in the existing literature and limitations of this review necessitate future studies and evidence synthesis to better understand DV and associated factors in diverse population group to strengthen the evidence base and inform better policies and practices.

Funding: No funding was received at any stage of conducting this review

Conflicts of interest: We declare no conflicts of interest

References

- Aromataris, E., Fernandez, R., Godfrey, C. M., Holly, C., Khalil, H., & Tungpunkom, P. (2015). Summarizing systematic reviews: methodological development, conduct and reporting of an umbrella review approach review of reviews, systematic review, umbrella review, umbrella review methodology. *International Journal of Evidence-Based Healthcare*.
<https://doi.org/10.1097/XEB.0000000000000055>
- Breiding, M. J., Basile, K. C., Smith, S. G., Black, M. C., & Mahendra, R. (2015). Intimate Partner Violence Surveillance: Uniform Definitions and Recommended Data Elements. *National Center for Injury Prevention and Control, Centers for Disease Control and Prevention*, 1–164. <https://doi.org/10.1136/bmj.328.7440.595>

- Burton, C. W., Halpern-Felsher, B., Rehm, R. S., Rankin, S. H., & Humphreys, J. C. (2016). Depression and Self-Rated Health Among Rural Women Who Experienced Adolescent Dating Abuse: A Mixed Methods Study. *Journal of Interpersonal Violence*, 31(5), 920–941. <https://doi.org/10.1177/0886260514556766>
- Caridade, S., Braga, T., & Borrajo, E. (2019). Cyber dating abuse (CDA): Evidence from a systematic review. *Aggression and Violent Behavior*, 48(May), 152–168. <https://doi.org/10.1016/j.avb.2019.08.018>
- Carlos, D. M., Campeiz, A. B., da Silva, J. L., Domingues Fernandes, M. I., da Cruz Leitão, M. N., Iossi Silv, M. A., & Carvalho Ferriani, M. das G. (2017). School-based interventions for teen dating violence prevention: integrative literature review. *Intervenciones En La Escuela Para Prevenir La Violencia En Las Relaciones Íntimas Entre Adolescentes: Revisión Integradora de La Literatura.*, 4(14), 133–145. Retrieved from <http://10.0.49.163/RIV17030>
- Center for Disease Control and Prevention (CDC). (2019). Preventing Teen Dating Violence. Retrieved October 22, 2019, from <https://www.cdc.gov/violenceprevention/intimatepartnerviolence/teendatingviolence/fastfact.html>
- Das Gupta, M., Engelman, R., Levy, J., Luchsinger, G., Merrick, T., Rosen, J. E., ... Adviser, R. (2014). *The State of World Population 2014 RESEARCHERS AND AUTHORS Other contributions from Sarah Castle UNFPA ADVISORY TEAM.*
- Draucker, C. B., Martsolf, D. S., Crane, S., McCord, A. L., Romero, L., & Al-Khattab, H. A. (2019). A feasibility test of an online intervention to prevention dating violence in emerging

adults. *Archives of Psychiatric Nursing*, 33(1), 37–45.

<https://doi.org/10.1016/j.apnu.2018.09.002>

Duerksen, K. N., & Woodin, E. M. (2019). Cyber Dating Abuse Victimization: Links With Psychosocial Functioning. *Journal of Interpersonal Violence*, 088626051987298.

<https://doi.org/10.1177/0886260519872982>

Duval, A., Lanning, B. A., & Patterson, M. S. (2018). A Systematic Review of Dating Violence Risk Factors Among Undergraduate College Students. *Trauma, Violence & Abuse*,

1524838018782207–1524838018782207. <https://doi.org/10.1177/1524838018782207>

Foshee, V. A., Karriker-Jaffe, K. J., Reyes, H. L. M. N., Ennett, S. T., Suchindran, C., Bauman,

K. E., & Benefield, T. S. (2008). What Accounts for Demographic Differences in

Trajectories of Adolescent Dating Violence? An Examination of Intrapersonal and

Contextual Mediators. *Journal of Adolescent Health*, 42(6), 596–604.

<https://doi.org/10.1016/j.jadohealth.2007.11.005>

Garthe, R. C., Sullivan, T. N., & McDaniel, M. A. (2017). A meta-analytic review of peer risk factors and adolescent dating violence. *Psychology of Violence*, 45–57. Retrieved from

<http://10.0.4.13/vio0000040>

Hébert, M., Daspe, M.-È., Lapierre, A., Godbout, N., Blais, M., Fernet, M., & Lavoie, F. (2019).

A Meta-Analysis of Risk and Protective Factors for Dating Violence Victimization: The

Role of Family and Peer Interpersonal Context. *Trauma, Violence & Abuse*, 20(4), 574–

590. Retrieved from <http://10.0.4.153/1524838017725336>

Hickman, L. J., Jaycox, L. H., & Aronoff, J. (2004). Dating Violence among Adolescents:

Prevalence, Gender Distribution, and Prevention Program Effectiveness. *Trauma, Violence*,

& Abuse, 5(2), 123–142. <https://doi.org/10.1177/1524838003262332>

Howard, D. E., Debnam, K. J., & Wang, M. Q. (2013). Ten-year trends in physical dating violence victimization among U.S. adolescent females. *The Journal of School Health, 83*(6), 389–399. <https://doi.org/10.1111/josh.12042>

Howard, D. E., Debnam, K. J., Wang, M. Q., & Gilchrist, B. (n.d.). 10-year trends in physical dating violence victimization among U.S. adolescent males. *International Quarterly of Community Health Education, 32*(4), 283–305. <https://doi.org/10.2190/IQ.32.4.c>

Johnson, R. M., LaValley, M., Schneider, K. E., Musci, R. J., Pettoruto, K., & Rothman, E. F. (2017). Marijuana use and physical dating violence among adolescents and emerging adults: A systematic review and meta-analysis. *Drug And Alcohol Dependence, 174*, 47–57. <https://doi.org/10.1016/j.drugalcdep.2017.01.012>

Johnson, R. M., Parker, E. M., Rinehart, J., Nail, J., & Rothman, E. F. (2015). Neighborhood Factors and Dating Violence Among Youth: A Systematic Review. *American Journal of Preventive Medicine, 49*(3), 458–466. <https://doi.org/10.1016/j.amepre.2015.05.020>

Joly, L. E., & Connolly, J. (2016). Dating violence among high-risk young women: A systematic review using quantitative and qualitative methods. *Behavioral Sciences, 6*(1). <https://doi.org/10.3390/bs6010007>

Jouriles, E. N., Garrido, E., Rosenfield, D., & McDonald, R. (2009). Experiences of psychological and physical aggression in adolescent romantic relationships: Links to psychological distress. *Child Abuse and Neglect, 33*(7), 451–460. <https://doi.org/10.1016/j.chiabu.2008.11.005>

- Liberati, A., Altman, D. G., Tetzlaff, J., Mulrow, C., Gotzsche, P. C., Ioannidis, J. P., ... Moher, D. (2009). The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *Bmj*, 339, b2700. <https://doi.org/10.1136/bmj.b2700>
- Mcleroy, K. R., Bibeau, D., Steckler, A., & Glanz, K. (1988). An Ecological Perspective on Health Promotion Programs. *Health Education & Behavior*, 15(4), 351–377. <https://doi.org/10.1177/109019818801500401>
- Munn, Z., Tufanaru, C., & Aromataris, E. (2014). JBI's systematic reviews: data extraction and synthesis. *The American Journal of Nursing*, 114(7), 49–54. <https://doi.org/10.1097/01.NAJ.0000451683.66447.89>
- O'Brien, K. M., Sauber, E. W., Kearney, M. S., Venaglia, R. B., & Lemay, E. P. (2019). Evaluating the Effectiveness of an Online Intervention to Educate College Students About Dating Violence and Bystander Responses. *Journal of Interpersonal Violence*. <https://doi.org/10.1177/0886260519829769>
- Park, S., & Kim, S.-H. (2018). The power of family and community factors in predicting dating violence: A meta-analysis. *Aggression & Violent Behavior*, 40, 19–28. <https://doi.org/10.1016/j.avb.2018.03.002>
- Park, S., & Kim, S.-H. (2019). Who Are the Victims and Who Are the Perpetrators in Dating Violence?: Sharing the Role of Victim and Perpetrator. *Trauma, Violence & Abuse*, 20(5), 1524838017730648. <https://doi.org/10.1177/1524838017730648>
- Park, Y., Mulford, C., & Blachman-Demner, D. (2018). The Acute and Chronic Impact of Adolescent Dating Violence. In *Adolescent Dating Violence* (pp. 53–83). Elsevier.

<https://doi.org/10.1016/b978-0-12-811797-2.00003-7>

- Pedersen, P., & Thomas, C. D. (1992). Prevalence and correlates of dating violence in a Canadian university sample. *Canadian Journal of Behavioural Science / Revue Canadienne Des Sciences Du Comportement*, 24(4), 490–501. <https://doi.org/10.1037/h0078756>
- Rothman, E. F., McNaughton Reyes, L., Johnson, R. M., & LaValley, M. (2012, January). Does the alcohol make them do it? Dating violence perpetration and drinking among youth. *Epidemiologic Reviews*. <https://doi.org/10.1093/epirev/mxr027>
- Rubio-Garay, F., López-González, M. A., Carrasco, M. Á., & Amor, P. J. (2017). The prevalence of dating violence: A systematic review. *Papeles Del Psicólogo*, 38(2), 135–147. <https://doi.org/10.23923/pap.psicol2017.2831>
- Spencer, C. M., Toews, M. L., Anders, K. M., & Emanuels, S. K. (2019). Risk Markers for Physical Teen Dating Violence Perpetration: A Meta-Analysis. *Trauma, Violence, & Abuse*, 152483801987570. <https://doi.org/10.1177/1524838019875700>
- Stonard, K. E., Bowen, E., Lawrence, T. R., & Price, S. A. (2014). The relevance of technology to the nature, prevalence and impact of Adolescent Dating Violence and Abuse: A research synthesis. *Aggression and Violent Behavior*, 19(4), 390–417. <https://doi.org/10.1016/j.avb.2014.06.005>
- Taquette, S. R., & Maia Monteiro, D. L. (2019). Causes and consequences of adolescent dating violence: a systematic review. *Journal of Injury & Violence Research*, 11(2), 137–147. <https://doi.org/10.5249/jivr.v11i2.1061>
- Teitelman, A. M., Ratcliffe, S. J., Dichter, M. E., & Sullivan, C. M. (2008). Recent and past

intimate partner abuse and HIV risk among young women. *JOGNN - Journal of Obstetric, Gynecologic, and Neonatal Nursing*, 37(2), 219–227. <https://doi.org/10.1111/j.1552-6909.2008.00231.x>

Temple, J. R., Choi, H. J., Elmquist, J. A., Hecht, M., Miller-Day, M., Stuart, G. L., ... Wolford-Clevenger, C. (2016). Psychological Abuse, Mental Health, and Acceptance of Dating Violence Among Adolescents. *Journal of Adolescent Health*, 59(2), 197–202. <https://doi.org/10.1016/j.jadohealth.2016.03.034>

The US Department of Justice. (n.d.). Dating Violence | OVW. Retrieved October 20, 2019, from <https://www.justice.gov/ovw/dating-violence>

Viejo, C., Monks, C. P., Sánchez, V., & Ortega-Ruiz, R. (2016). Physical Dating Violence in Spain and the United Kingdom and the Importance of Relationship Quality. *Journal of Interpersonal Violence*, 31(8), 1453–1475. <https://doi.org/10.1177/0886260514567963>

WHO. (2012). *Understanding and addressing violence against women*.

Wincentak, K., Connolly, J., & Card, N. (2017). Teen dating violence: A meta-analytic review of prevalence rates. *Psychology of Violence*, 7(2), 224–241. <https://doi.org/10.1037/a0040194>

Zych, I., Viejo, C., Vila, E., & Farrington, D. P. (2019). School Bullying and Dating Violence in Adolescents: A Systematic Review and Meta-Analysis. *Trauma, Violence & Abuse*, 1524838019854460–1524838019854460. <https://doi.org/10.1177/1524838019854460>

Appendix 1: Critical appraisal of the systematic reviews and meta-analyses

Author and year of publication	Is the review question clearly and explicitly stated?	Were the inclusion criteria appropriate for the review question?	Was the search strategy appropriate?	Were the sources and resources used to search for studies adequate?	Were the criteria for appraising studies appropriate?	Was critical appraisal conducted by two or more reviewers independently?	Were the methods used to combine studies appropriate?	Was the likelihood of publication bias assessed?	Were recommendations for policy and/or practice supported by the reported data?	Were the specific directives for new research appropriate?
(Rothman et al., 2012)	Yes	Yes	Yes	Yes	Not sure	Yes	Yes	Yes	N/A	Yes
(Stonard et al., 2014)	Yes	Yes	Yes	Yes	Yes	No	Yes	N/A	Yes	Yes
(Johnson et al., 2015)	Yes	Yes	Yes	Not sure	Yes	No	Yes	N/A	Yes	Yes
(Joly & Connolly, 2016)	Yes	Not sure	Yes	Yes	Not sure	No	Yes	No	N/A	Yes
(Rubio-Garay et al., 2017)	Yes	Not sure	Yes	Yes	Yes	No	Yes	No	Yes	No
(Garthe et al., 2017)	Yes	Yes	Yes	Yes	Yes	No	Yes	N/A	Yes	Yes
(Johnson et al., 2017)	Yes	Yes	Yes	Not sure	Yes	No	Yes	Yes	Yes	Yes
(Wincentak et al., 2017)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
(Duval et al., 2018)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	N/A	Yes	Yes
(S. Park & Kim, 2018)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	N/A

(Caridad e et al., 2019)	Yes	Yes	Yes	Yes	Yes	No	N/A	N/A	N/A	Yes
(Hébert et al., 2019)	Yes	Yes	Yes	Yes	Yes	No	Yes	Yes	Yes	Yes
(S. Park & Kim, 2019)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
(Spencer et al., 2019)	Yes	Yes	Yes	Yes	Not Sure	No	Yes	Yes	Yes	Yes
(Taquette & Maia Monteiro , 2019)	Yes	Yes	Not sure	No	N/A	Yes	Not sure	N/A	Yes	Yes
(Zych et al., 2019)	Yes	Yes	Yes	Yes	N/A	Yes	Yes	Yes	Yes	Yes